Carbohydrate Research, 215 (1991) c11–c13
Elsevier Science Publishers B.V., Amsterdam

CONTENTS

Special issue: A collection of papers on liquid chromatography

Foreword
K. B. Hicks (Philadelphia, PA, U.S.A.) vii

High-performance anion-exchange chromatography of sugars and sugar alcohols on quaternary ammonium resins under alkaline conditions
T. J. Paskach, P. J. Reilly (Ames, IA, U.S.A.), H.-P. Lieker, and K. Thielecke (Braunschweig, Germany) 1

H.p.l.c. of 4-nitrophenyl-α-malto-oligosaccharides
C. Niemann, W. Saenger, R. Nuck (Berlin, F.R.G.), and B. Pfannemüller (Freiburg, Germany) 15

High performance reversed-phase chromatographic mapping of 2-pyridylamino derivatives of xyloglucan oligosaccharides
Z. El Rassi, D. Tedford, J. An, and A. Mort (Stillwater, OK, U.S.A.) 25

Selectivity control in the anion-exchange chromatographic determination of saccharides in dairy products using pulsed amperometric detection
J. van Riel and C. Olieman (Ede, The Netherlands) 39

High-performance size-exclusion chromatography and ELISA detection of extracellular polysaccharides from Mucorales
G. A. de Ruiter, A. W. van der Lugt, A. G. J. Voragen and F. M. Rombouts (Wageningen, The Netherlands) and S. H. W., Notermans (Bilthoven, The Netherlands) 47

An h.p.l.c. method for determining chain-length distribution in some glycogens
N. W. H. Cheetham, N. Hansawek and P. Saecou (Kensington, Australia) 59

High-performance liquid chromatography of mono- and oligo-saccharides on a graphitized carbon column
K. Koizumi, Y. Okada, and M. Fukuda (Nishinomiya, Japan) 67

Isolation of oligogalacturonic acids in gram quantities by preparative h.p.l.c.
A. Hotchkiss, Jr., K. B. Hicks, L. W. Doner and P. L. Irwin (Philadelphia, U.S.A.) 81

Intrinsic viscosity and molecular weight of pectin components
M. L. Fishman, D. T. Gillispie, and S. M. Sondey (Philadelphia, U.S.A.) and Y. S. El-Atawy (Giza, Egypt) 91

Analysis of oligouronates by reversed-phase ion-pair h.p.l.c.: Role of the mobile phase
A. Heyraud and C. Leonard (Grenoble, France) 105

Determination of furosine in biomedical samples employing an improved hydrolysis and high-performance liquid chromatographic technique
W. T. Cefalu, A. Bell-Farrow, Z. Q. Wang, and S. Ralapati (Winston-Salem, NC, U.S.A.) 117

Characterization of five isomers of branched cyclomaltohepataose (β CD) having degree of polymerization (d.p.) = 9: Reinvestigation of three positional isomers of diglucosyl-β CD
K. Koizumi, T. Tanimoto, Y. Okada, N. Nakanishi, N. Kato (Nishinomiya, Japan), Y. Takagi and H. Hashimoto (Yokohama, Japan) 127

Kinetic analysis of pectate lyases by high-performance liquid chromatography
J. F. Preston III and J. D. Rice (Gainesville, FL, U.S.A.) 137

Kinetic comparisons of trimer-generating pectate and alginate lyases by reversed-phase ion-pair liquid chromatography
J. F. Preston III, J. D. Rice, M. C. Chow, and B. J. Brown (Gainesville, FL, U.S.A.) . . . 147

Optimization of waveforms for pulsed amperometric detection (p.a.d.) of carbohydrates following separation by liquid chromatography
W. R. LaCourse and D. C. Johnson (Ames, IA, U.S.A.) 159

High-performance anion-exchange chromatography with pulsed amperometric detection of linear and branched glucose oligosaccharides
R. N. Ammeraal, G. A. Delgado, F. L. Tenbarge, and R. B. Friedman (Hammond, IN, U.S.A.) 179

Capillary zone electrophoresis of reducing mono- and oligo-saccharides as the borate complexes of their 3-methyl-1-phenyl-2-pyrazolin-5-one derivatives
S. Honda, S. Suzuki, A. Nose, K. Yamamoto, and K. Kakehi (Higashiosaka, Japan) . . . 193

Notes

Chromatography of 1-deoxy-1-nitroalditols on a cation-exchange resin in the lanthanum form
M. Petrušová, L. Petruš, and E. Lattová (Bratislave, Czechoslavakia) 199

Analyses of α-linked disaccharides of D-glucose by high-performance liquid chromatography
M. Shiota (Tokyo, Japan) and S. Kobayashi (Ibaraki, Japan) 203

Separation of high-mannose isomers form yeast and mammalian sources using high-pH anion-exchange chromatography
R. R. Townsend (San Francisco, CA, U.S.A.), P. H. Atkinson (Bronx, NY, U.S.A.), and R. B. Trimble (Albany, NY, U.S.A.) 211

Problems encountered during the extraction, purification, and chromatography of pectic fragments, and some solutions to them
A. J. Mort, B. M. Moerschbacher, M. L. Pierce and N. O. Maness (Stillwater, OK, U.S.A.) 219

End of special issue papers

Molecular motion in glassy water–malto-oligosaccharide (maltodextrin) mixtures as studied by conventional and saturation-transfer spin-probe e.s.r. spectroscopy
M. J. G. W. Roozen, M. A. Hemminga and P. Walstra (Wageningen, Netherland) 229

The conformation of 1,6-anhydrolactose and its hexa-acetate in solution
A. Rivera-Sagredo and J. Jiménez-Barbero (Madrid, Spain) 239

A molecular modelling study of the distortion of α-cyclodextrin (cyclomaltohexaose) in complexes with guest molecules
S. P. van Helden, M. J. van Drooge, A. J. Claessens, A. C. A. Jansen, and L. H. M. Janssen (Utrecht, The Netherlands) 251

Synthesis of disaccharides with (1→4)-β glycosidic linkages related to the 4- and 6-sulfates and the 4,6-disulfates of chondroitin
M. Zsiška (Oldenburg, Germany) and B. Meyer (Athens, GA, U.S.A.) 261

Synthesis of β-D-GlcA-(1→3)-β-D-Gal disaccharides with 4- and 6-sulfate groups and 4,6-disulfate groups
M. Zsiška (Oldenburg, Germany) and B. Meyer (Athens, GA, U.S.A.) 279

Structure of an acidic glycan present in the lipopolysaccharide extract from the reference strain for *Serratia marcescens* serogroup O18
D. Oxley and S. G. Wilkinson (Hull, Gt. Britain) 293

Structural studies of the *Vibrio cholerae* O:3 O-antigen polysaccharide
T. A. Chowdhury, P.-E. Jansson, B. Lindberg, J. Lindberg, B. Gustafsson, and T. Holme (Stockholm, Sweden) 303

Distinguishing mammalian sialidases by inhibition kinetics with novel derivatives of 5-acetamido-2,6-anhydro-3,5-dideoxy-D-*glycero*-D-*galacto*-non-2-enonic acid, an unsaturated derivative of *N*-acetylneuraminic acid
T. G. Warner (Memphis, TN, U.S.A.), A. Louie (South San Francisco, CA, U.S.A.), M. Potier (Montreal, Canada), A. Ribeiro (Durham, NC, U.S.A.) 315

Structural analysis of the heptose/hexose region of the lipopolysaccharide from *Escherichia coli* K-12 strain W3100
O. Holst, U. Zähringer, H. Brade (Borstel, Germany), and A. Zamojski (Warsaw, Poland) 323

Notes
The synthesis and characterisation of 6-*O*-L-*glycero*-D-*manno*-heptopyranosyl-D-glucopyranose
Z. Pakulski, A. Zamojski (Warsaw, Poland), O. Holst, and U. Zähringer (Borstel, Germany) 337

Configurational assignment of *N*-acetylneuraminic acid and analogues *via* the vicinal C,H coupling constants
S. Prytulla, J. Lauterwein, M. Klessinger, and J. Thiem (Münster, Germany) 345

Determination of the activation energy and rate constants for the chair-to-chair interconversion of *cis*-inositol by n.m.r. spectroscopy
K. Dill and L. Huang (Clemson, SC, USA) 351

Complexing of sugars and sugar alcohols with metal ions: a comparative study by ion-exchange chromatography
M. M. Hämäläinen and H. Lönnberg (Turku, Finland) 357

Preliminary communication
Iodonium ion-assisted glycosylation of alkyl (aryl) 1-thioglycosides: regulation of stereoselectivity and reactivity
H. M. Zuurmond, S. C. van der Laan, G. A. van der Marel, and H. van Boom (Leiden, The Netherlands) c1

Author index c5

Subject index c7

Contents (Vol. 215) c11